arXiv:1405.1776v3 [astro-ph.CO] 12 Sep 2015

Approximate Metric for a Rotating Deformed Mass

Francisco Frutos-Alfaro, [1,2]*, Paulo Montero-Camacho[1],
Miguel Araya-Arguedas[1,2], Javier Bonatti-González[1,3]
[1]School of Physics, University of Costa Rica, San José, Costa Rica
[2]Space Research Center, University of Costa Rica, San José, Costa Rica
[3]Nuclear Research Center, University of Costa Rica, San José, Costa Rica

September 15, 2015

Abstract

A new Kerr-like metric with quadrupole moment is obtained by means of perturbing the Kerr spacetime. The form of this new metric is simple as the Kerr metric. By comparison with the exterior Hartle-Thorne metric, it is shown that it could be matched to an interior solution. This approximate metric may represent the spacetime of a real astrophysical object with any Kerr rotation parameter a and slightly deformed.

1 Introduction

In 1963, R. P. Kerr [26] proposed a metric that describes a massive rotating object. Since then, a huge amount of papers about the structure and astrophysical applications of this spacetime appeared. Now, it is widely believed that this metric does not represent the spacetime of an astrophysical rotating object. In 1967, Hernández [23] stated that *reasonable perfect fluid type solutions which might serve as source of the Kerr metric may not exist.* In

*E-mail: frutos@fisica.ucr.ac.cr

1971, Thorne [43, 32] added that *because of the relationship between multipole moments and angular momentum, the Kerr solution cannot represent correctly the external field of any realistic stars*. Moreover, the Kerr metric has difficulties when matching it to a realistic interior metric according to [3]. However, there has been a considerable amount of efforts trying to match the Kerr metric with a realistic interior metric that represents a physical source, see for example [7, 16, 17, 19, 28, 39]. For a concise and comprehensive review of the different methods that have been used in order to try and obtain an interior solution for the Kerr metric, see [27].

In [9] and [44], the Newman-Janis algorithm was applied to look for interior solutions. Drake and Turolla [9] also propose a general method for finding interior solutions with oblate spheroidal boundary surfaces and note that the boundary surfaces reduce to a sphere in the case with no rotation, however Vaggiu in [44] argues that it is more helpful to start with the Schwarzschild interior and then proceed to the Kerr interior. Vaggiu uses an anisotropic conformally flat static interior and is lead to interior Kerr solutions with oblate spheroidal boundary surfaces, additionally he points out that his procedure can be applied to find interior solutions matching with a general asymptotically flat vacuum stationary spacetime.

Other exact rotating solutions to the Einstein field equations (EFE) containing mass multipoles and magnetic dipole were obtained by [4, 30, 31, 33, 34, 35, 36, 37]. In the first four articles, they used the Ernst formalism [10], while in the four last ones, the solutions were obtained with the help of the Hoenselaers-Kinnersley-Xanthopoulos (HKX) transformations [25]. These authors obtain new metrics from a given seed metric. These formalisms allow to include other desirable characteristics (rotation, multipole moments, magnetic dipole, etc.) to a given seed metrics. Furthermore, Quevedo in [38] not only presents an exact electrovacuum solution that can be used to describe the exterior gravitational field of a rotating charged mass distribution, but also considers the matching using the derivatives of the curvature eigenvalues, this leads to matching conditions from which one can expect to obtain the minimum radius at which the matching can be made.

In Nature, it is expected that astrophysical objects are rotating and slightly deformed as is pointed out in [1] and in [42]. In addition, Andersson and Comer in [1] use a two fluid model for a neutron star one, layer with neutrons that has a differential rotation and another layer consisting of a solid crust with constant rotation. The aim of this article is to derive an appropriate analytical tractable metric for calculations in which the quadrupole

moment can be treated as perturbation, but for arbitrary angular momentum. Moreover, this metric should be useful to tackle astrophysical problems, for instance, accretion disk in compact stellar objects [1, 12, 21, 42], relativistic magnetohydrodynamic jet formation [11], astrometry [15, 41] and gravitational lensing [13]. Furthermore, software related with applications of the Kerr metric can be easily modified in order to include the quadrupole moment [8, 14, 45].
This paper is organized as follows. In section 2, we give a succinct explanation of the Kerr metric, and the weak limit of the Erez-Rosen metric is presented. In section 3, the Lewis metric is presented, and the perturbation method is discussed. The application of this method leads to a new solution to the EFE with quadrupole moment and rotation. It is checked by means of the REDUCE software [22] that the resulting metric is a solution of the EFE. In section 4, we compare our solution with the exterior Hartle-Thorne metric in order to assure that our metric has astrophysical meaning. Forthcoming works with this metric are discussed inx section 6.

2 The Kerr Metric and The Erez-Rosen Metric

2.1 The Kerr Metric

The Kerr metric represents the spacetime of a non-deformed massive rotating object. The Kerr metric is given by [26, 4]

$$\begin{aligned} ds^2 &= \frac{\Delta}{\rho^2}[dt - a\sin^2\theta d\phi]^2 - \frac{\sin^2\theta}{\rho^2}[(r^2+a^2)d\phi - adt]^2 \\ &- \frac{\rho^2}{\Delta}dr^2 - \rho^2 d\theta^2, \end{aligned} \tag{1}$$

where $\Delta = r^2 - 2Mr + a^2$ and $\rho^2 = r^2 + a^2\cos^2\theta$. M and a represent the mass and the rotation parameter, respectively. The angular momentum of the object is $J = Ma$.

2.2 The Erez-Rosen Metric

The Erez-Rosen metric [4, 46, 47, 48] represents the spacetime of a body with quadrupole moment. The principal axis of the quadrupole moment is chosen along the spin axis, so that gravitational radiation can be ignored. Here, we write down an approximate expression for this metric obtained by doing Taylor series [15]

$$\begin{aligned} ds^2 &= \left(1-\frac{2M}{r}\right)\mathrm{e}^{-2\chi}dt^2 - \left(1-\frac{2M}{r}\right)^{-1}\mathrm{e}^{2\chi}dr^2 \\ &- r^2\mathrm{e}^{2\chi}(d\theta^2+\sin^2\theta d\phi^2), \end{aligned} \tag{2}$$

where $d\Sigma^2 = d\theta^2 + \sin^2\theta d\phi^2$, and

$$\chi = \frac{2}{15}q\frac{M^3}{r^3}P_2(\cos\theta), \tag{3}$$

where $P_2(\cos\theta) = (3\cos^2\theta - 1)/2$. The quadrupole parameter is given by $q = 15GQ/(2c^2M^3)$, with Q representing the quadrupole moment. This metric is valid up to the order $O(qM^4, q^2)$.

3 Perturbing the Kerr Metric

3.1 The Lewis Metrics

The Lewis metric is given by [29, 4]

$$ds^2 = Vdt^2 - 2Wdtd\phi - \mathrm{e}^{\mu}d\rho^2 - \mathrm{e}^{\nu}dz^2 - Zd\phi^2, \tag{4}$$

where we have chosen the canonical coordinates $x^1 = \rho$ and $x^2 = z$, V, W, Z, μ and ν are functions of ρ and z ($\rho^2 = VZ + W^2$). Choosing $\mu = \nu$ and performing the following changes of potentials

$$V = f, \quad W = \omega f, \quad Z = \frac{\rho^2}{f} - \omega^2 f \quad \text{and} \quad \mathrm{e}^{\mu} = \frac{\mathrm{e}^{\gamma}}{f},$$

we get the Papapetrou metric

$$ds^2 = f(dt-\omega d\phi)^2 - \frac{\mathrm{e}^{\gamma}}{f}[d\rho^2 + dz^2] - \frac{\rho^2}{f}d\phi^2. \tag{5}$$

3.2 The Perturbation Method

To include a small quadrupole moment into the Kerr metric we will modify the Lewis-Papapetrou metric (5). First of all, we choose expressions for the canonical coordinates ρ and z. For the Kerr metric [26], one particular choice is [4, 6]

$$\rho = \sqrt{\Delta}\sin\theta \qquad \text{and} \qquad z = (r-M)\cos\theta, \tag{6}$$

where $\Delta = r^2 - 2Mr + a^2$.
From (6) we get

$$d\rho^2 + dz^2 = [(r-M)^2\sin^2\theta + \Delta\cos^2\theta]\left(\frac{dr^2}{\Delta} + d\theta^2\right). \tag{7}$$

If we choose

$$\mathrm{e}^{\mu} = \tilde{\rho}^2[(r-M)^2\sin^2\theta + \Delta\cos^2\theta]^{-1},$$

the term (7) becomes

$$e^{\mu}[d\rho^2 + dz^2] = \tilde{\rho}^2\left(\frac{dr^2}{\Delta} + d\theta^2\right),$$

where $\tilde{\rho}^2 = r^2 + a^2\cos^2\theta$.
From (5), we propose the following metric

$$ds^2 = \mathcal{V}dt^2 - 2\mathcal{W}dtd\phi - \mathcal{X}dr^2 - \mathcal{Y}d\theta^2 - \mathcal{Z}d\phi^2, \tag{8}$$

where

$$\begin{aligned} \mathcal{V} &= V\mathrm{e}^{-2\psi} \\ \mathcal{W} &= W \\ \mathcal{X} &= X\mathrm{e}^{2\psi} \\ \mathcal{Y} &= Y\mathrm{e}^{2\psi} \\ \mathcal{Z} &= Z\mathrm{e}^{2\psi}, \end{aligned} \tag{9}$$

where the potentials V, W, X, Y, Z, and ψ depend on $x^1 = r$ and $x^2 = \theta$. The potential $\mathcal{W} = W$ is so chosen to maintain the same cross components of the Kerr metric.

Now, let us choose

$$\begin{aligned} V &= f = \frac{1}{\tilde{\rho}^2}[\Delta - a^2 \sin^2\theta] \\ W &= \frac{a}{\tilde{\rho}^2}[\Delta - (r^2 + a^2)] \sin^2\theta = -\frac{2Jr}{\tilde{\rho}^2} \sin^2\theta \\ X &= \frac{\tilde{\rho}^2}{\Delta} \\ Y &= \tilde{\rho}^2 \\ Z &= \frac{\sin^2\theta}{\tilde{\rho}^2}[(r^2 + a^2)^2 - a^2 \Delta \sin^2\theta]. \end{aligned} \tag{10}$$

The only potential we have to find is ψ. In order to obtain this potential, the EFE must be solved

$$G_{ij} = R_{ij} - \frac{R}{2} g_{ij} = 0, \tag{11}$$

where R_{ij} (i, j = 0, 1, 2, 3) are the Ricci tensor components and R is the curvature scalar. The Ricci tensor components and the curvature scalar R for this metric can be found in the Appendix.
In our calculations, we consider the potential ψ as perturbation, *i.e.* one neglects terms of the form

$$\left(\frac{\partial\psi}{\partial r}\right)^2 = \left(\frac{\partial\psi}{\partial\theta}\right)^2 = \frac{\partial\psi}{\partial r}\frac{\partial\psi}{\partial\theta} \sim 0.$$

Terms containing factors of the form

$$a\frac{\partial\psi}{\partial x^i} = m\frac{\partial\psi}{\partial x^i} \sim 0 \qquad (i = 1, 2)$$

are also neglected. Substituting the known potentials (V, W, X, Y, Z) into the expressions for the Ricci tensor and the curvature scalar (see Appendix), it results only one equation for ψ that we have to solved:

$$r^2 \sin\theta \nabla^2\psi = \sin\theta\frac{\partial}{\partial r}\left(r^2\frac{\partial\psi}{\partial r}\right) + \frac{\partial}{\partial\theta}\left(\sin\theta\frac{\partial\psi}{\partial\theta}\right) = 0. \tag{12}$$

The solution for this equation is

$$\psi = \frac{\mathcal{K}}{r^3} P_2(\cos\theta), \tag{13}$$

where $\mathcal{K}$ is a constant. To determine this constant, we compare the weak limit of the metric (8) with the approximate Erez-Rosen metric (2). The result is $\mathcal{K} = 2qM^3/15$ ($\psi = \chi$).
Then, the new modified Kerr metric containing quadrupole moment is

$$\begin{aligned} ds^2 &= \frac{\mathrm{e}^{-2\chi}}{\rho^2}[\Delta - a^2\sin^2\theta]dt^2 + \frac{4Jr}{\rho^2}\sin^2\theta dt d\phi \\ &- \frac{\rho^2\mathrm{e}^{2\chi}}{\Delta}dr^2 - \rho^2\mathrm{e}^{2\chi}d\theta^2 \\ &- \frac{\mathrm{e}^{2\chi}\sin^2\theta}{\rho^2}[(r^2+a^2)^2 - a^2\Delta\sin^2\theta]d\phi^2 \\ &= \frac{\Delta}{\rho^2}[\mathrm{e}^{-\chi}dt - a\mathrm{e}^{\chi}\sin^2\theta d\phi]^2 \\ &- \frac{\sin^2\theta}{\rho^2}[(r^2+a^2)\mathrm{e}^{\chi}d\phi - a\mathrm{e}^{-\chi}dt]^2 \\ &- \mathrm{e}^{2\chi}\left(\frac{\rho^2}{\Delta}dr^2 + \rho^2 d\theta^2\right), \end{aligned} \tag{14}$$

where the tilde over ρ is dropped.
We verified that the metric (14) is indeed a solution of the EFE using REDUCE [22] up to the order $O(qM^4, q^2)$.
Note that (14) has four important limiting cases. One obtains the Kerr metric (1) if $q = 0$, the weak metric of [15] if $a^2 = q^2 = qM^4 \simeq 0$, the Erez-Rosen-like metric (2) if $a = 0$, and the Schwarzchild metric if $q = a = 0$.

4 Comparison with the Exterior Hartle-Thorne Metric

In order to validate the metric (14) as representing the gravitational field of a real astrophysical object, one should show that it is possible to construct an interior solution, which can appropriately be matched with our exterior solution. To this aim, we employed the exterior Hartle-Thorne metric [20, 2, 3, 15]

$$
\begin{aligned}
ds^2 &= \left(1-\frac{2\mathcal{M}}{r}+\frac{2\mathcal{Q}\mathcal{M}^3}{r^3}P_2(\cos\theta)\right)dt^2 \\
&- \left(1+\frac{2\mathcal{M}}{r}+\frac{4\mathcal{M}^2}{r^2}-\frac{2\mathcal{Q}\mathcal{M}^3}{r^3}P_2(\cos\theta)\right)dr^2 \\
&- r^2\left(1-\frac{2\mathcal{Q}\mathcal{M}^3}{r^3}P_2(\cos\theta)\right)d\Sigma^2+\frac{4\mathcal{J}}{r}\sin^2\theta dt d\phi,
\end{aligned}
\tag{15}
$$

where $\mathcal{M}$, $\mathcal{J}$, and $\mathcal{Q}$ are related with the total mass, angular momentum, and mass quadrupole moment of the rotating object, respectively. This approximation for the Hartle-Thorne metric (15) was obtained by Frutos-Alfaro *et al.* using a REDUCE program [15].
The spacetime (14) has the same weak limit as the metric obtained by Frutos-Alfaro *et al.* [15]. A comparison of the exterior Hartle-Thorne metric (15) with the weak limit of the metric (14) shows that upon defining

$$
\mathcal{M}=M, \qquad \mathcal{J}=J, \qquad 2\mathcal{Q}\mathcal{M}^3=-\frac{4}{15}qM^3, \tag{16}
$$

both metrics coincide up to the order $O(M^3,\, a^2,\, qM^4,\, q^2)$. Hence, the metric (14) may be used to represent a compact astrophysical object.

5 Conclusions

The new Kerr metric with quadrupole moment was obtained by solving the EFE approximately. It may represent the spacetime of a rotating and slightly deformed astrophysical object. This is possible, because it could be matched to an interior solution. We showed this by comparison of our metric with the exterior Hartle-Thorne metric. The limiting cases for the new Kerr metric correspond to the Kerr metric, the Erez-Rosen-like metric (see section 4), and the Schwarzschild metric as expected.
The inclusion of the quadrupole moment in the Kerr metric does it more suitable for astrophysical calculations than the Kerr metric alone. There are a large variety of applications which can be tackled with this new metric. Amongst the applications for this metric are astrometry, gravitational lensing, relativistic magnetohydrodynamic jet formation, and accretion disks in compact stellar objects, additionally we would like to point out that works in superfluid neutron stars can be repeated but using this new metric instead

of the Hartle-Thorne metric as an exterior solution. Furthermore, the existing software with applications of the Kerr metric can be easily modified to include the quadrupole moment.

A Appendix

The non-null Ricci tensor components for the metric (8) are given by (with the tilde over ρ dropped)

$$
\begin{aligned}
R_{00} &= \frac{\mathrm{e}^{-2\psi}}{4\rho^2X^2Y^2}\left(-4\rho^2VX^2Y\frac{\partial^2\psi}{\partial\theta^2}\right.\\
&+ 8VW^2X^2Y\left(\frac{\partial\psi}{\partial\theta}\right)^2 - 2\rho^2VXY\frac{\partial\psi}{\partial\theta}\frac{\partial X}{\partial\theta}\\
&+ 2VX^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial\rho^2}{\partial\theta} - 4\rho^2X^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial V}{\partial\theta}\\
&- 4W^2X^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial V}{\partial\theta} + 2\rho^2VX^2\frac{\partial\psi}{\partial\theta}\frac{\partial Y}{\partial\theta}\\
&- 4V^2X^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial Z}{\partial\theta} - 4\rho^2VXY^2\frac{\partial^2\psi}{\partial r^2}\\
&+ 8VW^2XY^2\left(\frac{\partial\psi}{\partial r}\right)^2 + 2\rho^2VY^2\frac{\partial\psi}{\partial r}\frac{\partial X}{\partial r}\\
&+ 2VXY^2\frac{\partial\psi}{\partial r}\frac{\partial\rho^2}{\partial r} - 4\rho^2XY^2\frac{\partial\psi}{\partial r}\frac{\partial V}{\partial r}\\
&- 4W^2XY^2\frac{\partial\psi}{\partial r}\frac{\partial V}{\partial r} - 2\rho^2VXY\frac{\partial\psi}{\partial r}\frac{\partial Y}{\partial r}\\
&- 4V^2XY^2\frac{\partial\psi}{\partial r}\frac{\partial Z}{\partial r} + \rho^2XY\frac{\partial X}{\partial\theta}\frac{\partial V}{\partial\theta}\\
&- \rho^2Y^2\frac{\partial X}{\partial r}\frac{\partial V}{\partial r} - X^2Y\frac{\partial\rho^2}{\partial\theta}\frac{\partial V}{\partial\theta}\\
&- XY^2\frac{\partial\rho^2}{\partial r}\frac{\partial V}{\partial r} + 2\rho^2X^2Y\frac{\partial^2V}{\partial\theta^2}\\
&- \rho^2X^2\frac{\partial V}{\partial\theta}\frac{\partial Y}{\partial\theta} + 2VX^2Y\frac{\partial V}{\partial\theta}\frac{\partial Z}{\partial\theta}\\
&+ 2\rho^2XY^2\frac{\partial^2V}{\partial r^2} + \rho^2XY\frac{\partial V}{\partial r}\frac{\partial Y}{\partial r}\\
&+ 2VXY^2\frac{\partial V}{\partial r}\frac{\partial Z}{\partial r} + 2VX^2Y\left(\frac{\partial W}{\partial\theta}\right)^2\\
&+ \left.2VXY^2\left(\frac{\partial W}{\partial r}\right)^2\right)
\end{aligned}
$$

$$
\begin{aligned}
R_{03} &= \frac{\mathrm{e}^{-2\psi}}{4\rho^2X^2Y^2}\left(8\rho^2WX^2Y\left(\frac{\partial\psi}{\partial\theta}\right)^2\right.\\
&- 8W^3X^2Y\left(\frac{\partial\psi}{\partial\theta}\right)^2 - 4WX^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial\rho^2}{\partial\theta}\\
&+ 8W^2X^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial W}{\partial\theta} + 8VWX^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial Z}{\partial\theta}\\
&+ 8\rho^2WXY^2\left(\frac{\partial\psi}{\partial r}\right)^2 - 8W^3XY^2\left(\frac{\partial\psi}{\partial r}\right)^2\\
&- 4WXY^2\frac{\partial\psi}{\partial r}\frac{\partial\rho^2}{\partial r} + 8W^2XY^2\frac{\partial\psi}{\partial r}\frac{\partial W}{\partial r}\\
&+ 8VWXY^2\frac{\partial\psi}{\partial r}\frac{\partial Z}{\partial r} - \rho^2XY\frac{\partial X}{\partial\theta}\frac{\partial W}{\partial\theta}\\
&+ \rho^2Y^2\frac{\partial X}{\partial r}\frac{\partial W}{\partial r} + X^2Y\frac{\partial\rho^2}{\partial\theta}\frac{\partial W}{\partial\theta}\\
&+ XY^2\frac{\partial\rho^2}{\partial r}\frac{\partial W}{\partial r} - 2WX^2Y\frac{\partial V}{\partial\theta}\frac{\partial Z}{\partial\theta}\\
&- 2WXY^2\frac{\partial V}{\partial r}\frac{\partial Z}{\partial r} - 2\rho^2X^2Y\frac{\partial^2W}{\partial\theta^2}\\
&- 2WX^2Y\left(\frac{\partial W}{\partial\theta}\right)^2 + \rho^2X^2\frac{\partial W}{\partial\theta}\frac{\partial Y}{\partial\theta}\\
&- 2\rho^2XY^2\frac{\partial^2W}{\partial r^2} - 2WXY^2\left(\frac{\partial W}{\partial r}\right)^2\\
&- \left.\rho^2XY\frac{\partial W}{\partial r}\frac{\partial Y}{\partial r}\right) = R_{30}
\end{aligned}
$$

$$
\begin{aligned}
R_{11} &= \frac{1}{4\rho^4 XY^2}\left(-4\rho^4X^2Y\frac{\partial^2\psi}{\partial\theta^2} - 2\rho^4XY\frac{\partial\psi}{\partial\theta}\frac{\partial X}{\partial\theta}\right.\\
&- 2\rho^2X^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial\rho^2}{\partial\theta} + 2\rho^4X^2\frac{\partial\psi}{\partial\theta}\frac{\partial Y}{\partial\theta}\\
&- 4\rho^4XY^2\frac{\partial^2\psi}{\partial r^2} - 8\rho^4XY^2\left(\frac{\partial\psi}{\partial r}\right)^2\\
&+ 8\rho^2W^2XY^2\left(\frac{\partial\psi}{\partial r}\right)^2 + 2\rho^4Y^2\frac{\partial\psi}{\partial r}\frac{\partial X}{\partial r}\\
&+ 6\rho^2XY^2\frac{\partial\psi}{\partial r}\frac{\partial\rho^2}{\partial r} - 8\rho^2WXY^2\frac{\partial\psi}{\partial r}\frac{\partial W}{\partial r}\\
&- 2\rho^4XY\frac{\partial\psi}{\partial r}\frac{\partial Y}{\partial r} - 8\rho^2VXY^2\frac{\partial\psi}{\partial r}\frac{\partial Z}{\partial r}\\
&- 2\rho^4XY\frac{\partial^2X}{\partial\theta^2} + \rho^4Y\left(\frac{\partial X}{\partial\theta}\right)^2\\
&- \rho^2XY\frac{\partial X}{\partial\theta}\frac{\partial\rho^2}{\partial\theta} + \rho^4X\frac{\partial X}{\partial\theta}\frac{\partial Y}{\partial\theta}\\
&+ \rho^2Y^2\frac{\partial X}{\partial r}\frac{\partial\rho^2}{\partial r} + \rho^4Y\frac{\partial X}{\partial r}\frac{\partial Y}{\partial r}\\
&- 2\rho^2XY^2\frac{\partial^2\rho^2}{\partial r^2} + XY^2\left(\frac{\partial\rho^2}{\partial r}\right)^2\\
&+ 2VXY^2\frac{\partial\rho^2}{\partial r}\frac{\partial Z}{\partial r} + 2W^2XY^2\frac{\partial V}{\partial r}\frac{\partial Z}{\partial r}\\
&+ 2\rho^2XY^2\left(\frac{\partial W}{\partial r}\right)^2 - 4VWXY^2\frac{\partial W}{\partial r}\frac{\partial Z}{\partial r}\\
&- 2\rho^4XY\frac{\partial^2Y}{\partial r^2} + \rho^4X\left(\frac{\partial Y}{\partial r}\right)^2\\
&\left.- 2V^2XY^2\left(\frac{\partial Z}{\partial r}\right)^2\right)
\end{aligned}
$$

$$
\begin{aligned}
R_{12} &= \frac{1}{4\rho^4 XY}\left(-8\rho^4 XY\frac{\partial\psi}{\partial\theta}\frac{\partial\psi}{\partial r}\right.\\
&+ 8\rho^2W^2XY\frac{\partial\psi}{\partial\theta}\frac{\partial\psi}{\partial r} + 4\rho^2XY\frac{\partial\psi}{\partial\theta}\frac{\partial\rho^2}{\partial r}\\
&- 4\rho^2WXY\frac{\partial\psi}{\partial\theta}\frac{\partial W}{\partial r} - 4\rho^2VXY\frac{\partial\psi}{\partial\theta}\frac{\partial Z}{\partial r}\\
&+ 4\rho^2XY\frac{\partial\psi}{\partial r}\frac{\partial\rho^2}{\partial\theta} - 4\rho^2WXY\frac{\partial\psi}{\partial r}\frac{\partial W}{\partial\theta}\\
&- 4\rho^2VXY\frac{\partial\psi}{\partial r}\frac{\partial Z}{\partial\theta} + \rho^2Y\frac{\partial X}{\partial\theta}\frac{\partial\rho^2}{\partial r}\\
&- 2\rho^2XY\frac{\partial^2\rho^2}{\partial\theta\partial r} + W^2XY\frac{\partial^2\rho^2}{\partial\theta\partial r}\\
&+ XY\frac{\partial\rho^2}{\partial\theta}\frac{\partial\rho^2}{\partial r} + \rho^2X\frac{\partial\rho^2}{\partial\theta}\frac{\partial Y}{\partial r}\\
&+ VXY\frac{\partial\rho^2}{\partial\theta}\frac{\partial Z}{\partial r} + VXY\frac{\partial\rho^2}{\partial r}\frac{\partial Z}{\partial\theta}\\
&- W^2XYZ\frac{\partial^2V}{\partial\theta\partial r} - 2W^3XY\frac{\partial^2W}{\partial\theta\partial r}\\
&+ 2\rho^2XY\frac{\partial W}{\partial\theta}\frac{\partial W}{\partial r} - 2W^2XY\frac{\partial W}{\partial\theta}\frac{\partial W}{\partial r}\\
&- 2VWXY\frac{\partial W}{\partial\theta}\frac{\partial Z}{\partial r} - 2VWXY\frac{\partial W}{\partial r}\frac{\partial Z}{\partial\theta}\\
&- \left.VW^2XY\frac{\partial^2Z}{\partial\theta\partial r} - 2V^2XY\frac{\partial Z}{\partial\theta}\frac{\partial Z}{\partial r}\right) = R_{21}
\end{aligned}
$$

$$
\begin{aligned}
R_{22} &= \frac{1}{4\rho^4 X^2 Y}\left(-4\rho^4 X^2 Y \frac{\partial^2\psi}{\partial\theta^2}\right. \\
&- 8\rho^4 X^2 Y\left(\frac{\partial\psi}{\partial\theta}\right)^2 + 8\rho^2 W^2 X^2 Y\left(\frac{\partial\psi}{\partial\theta}\right)^2 \\
&- 2\rho^4 XY\frac{\partial\psi}{\partial\theta}\frac{\partial X}{\partial\theta} + 6\rho^2 X^2 Y\frac{\partial\psi}{\partial\theta}\frac{\partial\rho^2}{\partial\theta} \\
&- 8\rho^2 W X^2 Y\frac{\partial\psi}{\partial\theta}\frac{\partial W}{\partial\theta} + 2\rho^4 X^2\frac{\partial\psi}{\partial\theta}\frac{\partial Y}{\partial\theta} \\
&- 8\rho^2 V X^2 Y\frac{\partial\psi}{\partial\theta}\frac{\partial Z}{\partial\theta} - 4\rho^4 XY^2\frac{\partial^2\psi}{\partial r^2} \\
&+ 2\rho^4 Y^2\frac{\partial\psi}{\partial r}\frac{\partial X}{\partial r} - 2\rho^2 XY^2\frac{\partial\psi}{\partial r}\frac{\partial\rho^2}{\partial r} \\
&- 2\rho^4 XY\frac{\partial\psi}{\partial r}\frac{\partial Y}{\partial r} - 2\rho^4 XY\frac{\partial^2 X}{\partial\theta^2} \\
&+ \rho^4 Y\left(\frac{\partial X}{\partial\theta}\right)^2 + \rho^4 X\frac{\partial X}{\partial\theta}\frac{\partial Y}{\partial\theta} \\
&+ \rho^4 Y\frac{\partial X}{\partial r}\frac{\partial Y}{\partial r} - 2\rho^2 X^2 Y\frac{\partial^2\rho^2}{\partial\theta^2} \\
&+ X^2 Y\left(\frac{\partial\rho^2}{\partial\theta}\right)^2 + \rho^2 X^2\frac{\partial\rho^2}{\partial\theta}\frac{\partial Y}{\partial\theta} \\
&+ 2V X^2 Y\frac{\partial\rho^2}{\partial\theta}\frac{\partial Z}{\partial\theta} - \rho^2 XY\frac{\partial\rho^2}{\partial r}\frac{\partial Y}{\partial r} \\
&+ 2W^2 X^2 Y\frac{\partial V}{\partial\theta}\frac{\partial Z}{\partial\theta} + 2\rho^2 X^2 Y\left(\frac{\partial W}{\partial\theta}\right)^2 \\
&- 4VW X^2 Y\frac{\partial W}{\partial\theta}\frac{\partial Z}{\partial\theta} - 2\rho^4 XY\frac{\partial^2 Y}{\partial r^2} \\
&+ \left.\rho^4 X\left(\frac{\partial Y}{\partial r}\right)^2 - 2V^2 X^2 Y\left(\frac{\partial Z}{\partial\theta}\right)^2\right)
\end{aligned}
$$

$$
\begin{aligned}
R_{33} &= \frac{1}{4\rho^2X^2Y^2}\left(-4\rho^2X^2YZ\frac{\partial^2\psi}{\partial\theta^2}\right.\\
&- 8W^2X^2YZ\left(\frac{\partial\psi}{\partial\theta}\right)^2 - 2\rho^2XYZ\frac{\partial\psi}{\partial\theta}\frac{\partial X}{\partial\theta}\\
&- 2YX^2Z\frac{\partial\psi}{\partial\theta}\frac{\partial\rho^2}{\partial\theta} + 8WX^2YZ\frac{\partial\psi}{\partial\theta}\frac{\partial W}{\partial\theta}\\
&+ 2\rho^2X^2Z\frac{\partial\psi}{\partial\theta}\frac{\partial Y}{\partial\theta} - 8W^2X^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial Z}{\partial\theta}\\
&- 4\rho^2XY^2Z\frac{\partial^2\psi}{\partial r^2} - 8W^2XY^2Z\left(\frac{\partial\psi}{\partial r}\right)^2\\
&+ 2\rho^2Y^2Z\frac{\partial\psi}{\partial r}\frac{\partial X}{\partial r} - 2XY^2Z\frac{\partial\psi}{\partial r}\frac{\partial\rho^2}{\partial r}\\
&+ 8WXY^2Z\frac{\partial\psi}{\partial r}\frac{\partial W}{\partial r} - 2\rho^2XYZ\frac{\partial\psi}{\partial r}\frac{\partial Y}{\partial r}\\
&- 8W^2XY^2\frac{\partial\psi}{\partial r}\frac{\partial Z}{\partial r} - \rho^2XY\frac{\partial X}{\partial\theta}\frac{\partial Z}{\partial\theta}\\
&+ \rho^2Y^2\frac{\partial X}{\partial r}\frac{\partial Z}{\partial r} - X^2Y\frac{\partial\rho^2}{\partial\theta}\frac{\partial Z}{\partial\theta}\\
&- XY^2\frac{\partial\rho^2}{\partial r}\frac{\partial Z}{\partial r} - 2X^2YZ\left(\frac{\partial W}{\partial\theta}\right)^2\\
&+ 4WX^2Y\frac{\partial W}{\partial\theta}\frac{\partial Z}{\partial\theta} - 2XY^2Z\left(\frac{\partial W}{\partial r}\right)^2\\
&+ 4WXY^2\frac{\partial W}{\partial r}\frac{\partial Z}{\partial r} + \rho^2X^2\frac{\partial Y}{\partial\theta}\frac{\partial Z}{\partial\theta}\\
&- \rho^2XY\frac{\partial Y}{\partial r}\frac{\partial Z}{\partial r} - 2\rho^2X^2Y\frac{\partial^2 Z}{\partial\theta^2}\\
&+ 2VX^2Y\left(\frac{\partial Z}{\partial\theta}\right)^2 - 2\rho^2XY^2\frac{\partial^2 Z}{\partial r^2}\\
&+ \left.2VXY^2\left(\frac{\partial Z}{\partial r}\right)^2\right)
\end{aligned}
$$

Calculation of the scalar curvature (with the tilde over ρ dropped)

$$
\begin{aligned}
R &= \frac{\mathrm{e}^{-2\psi}}{2\rho^4X^2Y^2}\left(4\rho^4X^2Y\frac{\partial^2\psi}{\partial\theta^2}+4\rho^4X^2Y\left(\frac{\partial\psi}{\partial\theta}\right)^2\right.\\
&- 4\rho^2W^2X^2Y\left(\frac{\partial\psi}{\partial\theta}\right)^2+2\rho^4XY\frac{\partial\psi}{\partial\theta}\frac{\partial X}{\partial\theta}\\
&- 2\rho^2X^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial\rho^2}{\partial\theta}+4\rho^2WX^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial W}{\partial\theta}\\
&- 2\rho^4X^2\frac{\partial\psi}{\partial\theta}\frac{\partial Y}{\partial\theta}+4\rho^2VX^2Y\frac{\partial\psi}{\partial\theta}\frac{\partial Z}{\partial\theta}\\
&+ 4\rho^4XY^2\frac{\partial^2\psi}{\partial r^2}+4\rho^4XY^2\left(\frac{\partial\psi}{\partial r}\right)^2\\
&- 4\rho^2W^2XY^2\left(\frac{\partial\psi}{\partial r}\right)^2-2\rho^4Y^2\frac{\partial\psi}{\partial r}\frac{\partial X}{\partial r}\\
&- 2\rho^2XY^2\frac{\partial\psi}{\partial r}\frac{\partial\rho^2}{\partial r}+4\rho^2WXY^2\frac{\partial\psi}{\partial r}\frac{\partial W}{\partial r}\\
&+ 2\rho^4XY\frac{\partial\psi}{\partial r}\frac{\partial Y}{\partial r}+4\rho^2VXY^2\frac{\partial\psi}{\partial r}\frac{\partial Z}{\partial r}\\
&+ 2\rho^4XY\frac{\partial^2X}{\partial\theta^2}-\rho^4Y\left(\frac{\partial X}{\partial\theta}\right)^2\\
&+ \rho^2XY\frac{\partial X}{\partial\theta}\frac{\partial\rho^2}{\partial\theta}-\rho^4X\frac{\partial X}{\partial\theta}\frac{\partial Y}{\partial\theta}\\
&- \rho^2Y^2\frac{\partial X}{\partial r}\frac{\partial\rho^2}{\partial r}-\rho^4Y\frac{\partial X}{\partial r}\frac{\partial Y}{\partial r}\\
&+ 2\rho^2X^2Y\frac{\partial^2\rho^2}{\partial\theta^2}-X^2Y\left(\frac{\partial\rho^2}{\partial\theta}\right)^2\\
&- \rho^2X^2\frac{\partial\rho^2}{\partial\theta}\frac{\partial Y}{\partial\theta}+2\rho^2XY^2\frac{\partial^2\rho^2}{\partial r^2}\\
&- XY^2\left(\frac{\partial\rho^2}{\partial r}\right)^2+\rho^2XY\frac{\partial\rho^2}{\partial r}\frac{\partial Y}{\partial r}\\
&- \rho^2X^2Y\frac{\partial V}{\partial\theta}\frac{\partial Z}{\partial\theta}-\rho^2XY^2\frac{\partial V}{\partial r}\frac{\partial Z}{\partial r}\\
&- \rho^2X^2Y\left(\frac{\partial W}{\partial\theta}\right)^2-\rho^2XY^2\left(\frac{\partial W}{\partial r}\right)^2\\
&+ \left.2\rho^4XY\frac{\partial^2Y}{\partial r^2}-\rho^4X\left(\frac{\partial Y}{\partial r}\right)^2\right)
\end{aligned}
$$

References

[1] Andersson, N., Comer, G. L., 2001, Class. Quantum Grav., 18, 969

[2] Berti, E., White, F., Maniopoulou, A., Bruni, M., 2005, MNRAS, 358, 923

[3] Boshkayev, K., Quevedo, H., Ruffini, R., 2012, PRD, 86, 064043

[4] Carmeli, M., 2001, Classical Fields, World Scientific Publishing, Singapore

[4] Castejon-Amenedo, J., Manko, V. S., 1990, PRD, 41, 2018

[6] Chandrasekhar, S., 2000, The Mathematical Theory of Black Holes, Oxford University Press, Oxford

[7] Cuchí, J. E., Molina, A., Ruiz, E., 2011, J. Phys.: Conf. Ser., 314, 012070

[8] Dexter, J., Algol, E., 2009, ApJ, 696, 1616

[9] Drake, S. P., Turolla, R., 1997, Class. Quantum Grav., 14, 1883

[10] Ernst, F. J., 1968, Phys. Rev., 167, 1175

[11] Fendt, C., Memola, E., 2008, Inter. Jour. of Modern Phys., D17, 1677

[12] Fragile, P. C., Blaes, O. M., Anninos, P., Salmonson, J. D., 2007, ApJ, 668, 417

[13] Frutos-Alfaro, F., 2001, Am. J. Phys., 69, 218

[14] Frutos-Alfaro, F., Grave, F., Müller, T., Adis, D., 2012, Jour. of Modern Phys., 3, 1882

[15] Frutos-Alfaro, F., Retana-Montenegro, E., Cordero-García, I., Bonatti-González, J., 2013, International Journal of Astronomy and Astrophysics, 3, 431 (ArXiv: 1209.6126v2)

[16] Glass, E. N., Krisch, J. P., 2009, Class. Quantum Grav., 26, 175010

[17] Haggag, S., Marek, J., 1981, Il Nuovo Cimento, 62B, 273

[18] Haggag, S., 1990, Il Nuovo Cimento, 105B, 365

[19] Haggag, S., 1990, Astrophys. and Space Sci., 173, 47

[20] Hartle, J. B., Thorne, K. S., 1968, ApJ, 153, 807

[21] Hawley, J. F., 2009, Astrophys. and Space Sci., 320, 107

[22] Hearn, A. C., 1999, REDUCE (User's and Contributed Packages Manual), Konrad-Zuse-Zentrum für Informationstechnik, Berlin

[23] Hernández, W., 1967, Physical Rev., 159, 1070.

[24] Hernández-Pastora, J. L., 2006, Gen. Rel. Grav., 38(5), 871.

[25] Hoenselaers, C., Kinnersley, W., Xanthopoulos, B. C., 1979, J. Math. Phys., 20(12), 2530

[26] Kerr, R. P., 1963, PRL, 11, 237

[27] Krasiński, A., 1978, Annals of Physics, 112, 22

[28] Krasiński, A., 1980, Physics Letters, 80A, 238

[29] Lewis. T., 1932, Proc. Roy. Soc. Lond., A, 176

[30] Manko, V. S., Novikov, I. D., 1992, Class. Quantum Grav., 9, 2477

[31] Manko, V. S., Mielke, E. W., Sanabria-Gómez, J. D., 2000, PRD, 61, 081501

[32] Marsh, G. E. (ArXiv: 1404.5297v3)

[33] Pachón, L. A., Rueda, J. A., Sanabria-Gómez, J. D., 2006, PRD, 73, 104038

[34] Quevedo, H., 1986, PRD, 33, 324

[35] Quevedo, H., 1989, PRD, 39, 2904

[36] Quevedo, H., Mashhoon, B., 1991, PRD, 43, 3902

[37] Quevedo, H., 2011, Gen. Rel. Grav., 43, 1141

[38] Quevedo, H., 2012, Proceedings of the Twelfth Marcel Grossmann Meeting on General Relativity, edited by Thibault Damour, Robert T. Jantzen and Remo Ruffini, World Scientific, Singapore (ArXiv: 1205.0500v1)

[39] Ramadan, A., 2004, Il Nuovo Cimento, 119B, 123

[40] Sato, H., Tomimatsu, A., 1973, Prog. Theor. Physics, 49(3), 790.

[41] Soffel, M. H., 1989, Relativity in Astrometry, Celestial Mechanics and Geodesy (Astronomy and Astrophysics Library), Springer-Verlag, Berlin

[42] Stergioulas, N., 2003, Living Reviews in Relativity, 6, 3

[43] Thorne, K. S., 1969, Relativistic Stars, Black Holes and Gravitational Waves in B. K. Sachs, ed., General Relativity and Cosmology, Proceedings of the International School of Physics Enrico Fermi, Course XLVII, pp. 237-283, Academic Press.

[44] Viaggiu, S., 2006, Int. J. Mod. Phys. D., 15, 1441

[45] Vincent, F. H., Paumard, T., Gourgoulhon, E., Perrin, G., 2011, Class. Quantum Grav., 28, 225011 (ArXiv: 1109.4769)

[46] Winicour, J., Janis, A. I., Newman, E. T., 1968, Phys. Rev., 176, 1507

[47] Young, J. H., Coulter, C. A., 1969, Phys. Rev., 184, 1313

[48] Zel'dovich, Ya. B., Novikov, I. D., 2011, Stars and Relativity, Dover Publications, New York